Morphos Modeling of Nearshore Morphology and Data Needs


Bradley Johnson Coastal and Hyd. Lab Vicksburg, MS

- Overall approach
- The spectrum of efforts
- Dutch approach and results (Xbeach)
- CHL/U. Delaware
 - CSHORE approach and results
 - •C2SHORE results
- Data needs to drive modeling



Approach

- Use open-water surge/current and wave models or data to drive nearshore FD model
- Nearshore model domain: depth of morphology change through dune and beyond
- Various approaches/levels of modeling to assure we have the appropriate set of tools to address project goals







Spectrum of Efforts

CSHORE CZSHORE

Theach

Unesco-IHE,

Phase resolving

-arge Computational effort

·CSHORE: profile development

·C2SHORE: 2DH quasi-steady

*Xbeach: low frequency resolving



XBeach


Approach

- Compiled code in modules
- Short-wave averaged but long-wave resolving modeling of waves, flow and morphology change in time-domain
- Swash and overwash motions
- Dune erosion, overwashing, breaching and full inundation
- Domain from outside surfzone to backbarrier
- Driven by boundary conditions from surge and spectral wave models



XBeach

Principle sketch - physics





XBeach


Sediment transport

- Depth-integrated advection-diffusion equation
- Equilibrium concentration determined by Soulsby-van Rijn formulations
- Velocity includes seaward return flow

$$\frac{\partial hC}{\partial t} + \frac{\partial hCu^{E}}{\partial x} + \frac{\partial hCv^{E}}{\partial y} + \frac{\partial}{\partial x} \left[D_{h}h \frac{\partial C}{\partial x} \right] + \frac{\partial}{\partial y} \left[D_{h}h \frac{\partial C}{\partial y} \right] = \frac{hC_{eq} - hC}{T_{s}}$$



XBeach Detailed beach process model plus avalanching





Approach

- Simple, Efficient, Wide Range of Verification
- Longshore Uniform Formulation
- Steady Formulation
- Shallow Water Hydrodynamics
- Probabalistic Representation of Sediment Transport
- Entrainment driven by Energy Dissipation
- Includes Wave and Current Transport
- Bed load and Suspended load



Suspended Sediment

• Suspended sediment volume V_s per unit area is estimated as

$$V_{s} = \frac{e_{B}D_{B} + e_{f}D_{f}}{\rho g(s-1)w_{f}}P_{s}$$


- where $e_B = 0.005$ and $e_f = 0.01$.
- Probability P_s is estimated as the probability of u' exceeding sediment fall velocity








Typical Application (East Coast)






Typical Application (East Coast)






Typical Application (West Coast)






Typical Application (West Coast)






OSU





OSU






Extension of CSHORE to 2DH

<u>STWAVE</u> + <u>STCIRC</u> + <u>Sediment Transport Model</u>


- Extension to 2DH Allows longshore variations, inlets, etc.
- Coupled wave, circulation, sediment transport, morphology



FRF During Isabel (model result)




Model Framework




Transport algorithms are and extension, but framework is more flexible.








Model domain






Model Run






Post Measured, Computed Bed





C2SHORE Post and Pre Difference





What does this reveal about C2SHORE?

- Water levels? --- two m of discrepancy between basin-scale models
- Waves? --- no data
- Currents --- no data
- Bathymetry --- only pre-storm



What does this reveal about C2SHORE?


- Water levels? --- two m of discrepancy between basin-scale models
- Waves? --- no data
- Currents --- no data
- Bathymetry --- only pre-storm

Revealed: C2SHORE is predicting morhpology change with some error, but why?



Waves Currents

at boundar Offshore waves




Water Levels

Geometry



Waves

Currents



Water Levels


Offshore water leve

Geometry



Waves

Currents



Water Levels

Geometry



Waves

Currents

- Sediment and underlying features
- Pre-storm conditions (topo + bathy + sed) at last minute
- Post-storm profiles shortly following



Water Levels

Site & Geometry



A Joint Effort

- A Willapa Bay model benefits District
 - A high resolution circulation Bay model
 - Detailed nearshore domain for Morphology
 - Some confidence with present tools
- Monitoring data benefits MORPHOS
 - High quality data of large events is needed
 - Complex system drives the technology

